



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

68Ga-PSMA PET/MR-positive peritoneal metastasis in the falciform ligament in recurrent prostate cancer

Kranzbühler, Benedikt ; Tran, Sébastien ; Zilli, Thomas ; Burger, Irene A

Abstract: Synchronous or metachronous metastases develop in approximately one third of all prostate cancer (PCa) patients. Main sites of metastasis include lymph nodes, bone, lung and liver. Secondary peritoneal carcinomatosis is very rare in PCa, with only a few published cases. Ga-PSMA PET-based imaging is a promising tool for staging and restaging PCa. We report a case of Ga-PSMA PET/MR-positive peritoneal metastasis as site of primary relapse after definitive PCa treatment in a 58-year-old man with a prostate-specific antigen of 30 ng/mL at time of the study. Exploratory laparoscopy and subsequent histopathologic examination confirmed nonascitic peritoneal PCa carcinomatosis.

DOI: <https://doi.org/10.1097/RLU.0000000000001703>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-137890>

Journal Article

Published Version

Originally published at:

Kranzbühler, Benedikt; Tran, Sébastien; Zilli, Thomas; Burger, Irene A (2017). 68Ga-PSMA PET/MR-positive peritoneal metastasis in the falciform ligament in recurrent prostate cancer. *Clinical Nuclear Medicine*, 42(8):e388-e389.

DOI: <https://doi.org/10.1097/RLU.0000000000001703>

⁶⁸Ga-PSMA PET/MR–Positive Peritoneal Metastasis in the Falciform Ligament in Recurrent Prostate Cancer

Benedikt Kranzbühler, MD,* Sébastien Tran, MD,† Thomas Zilli, MD,‡ and Irene A. Burger, MD‡

Abstract: Synchronous or metachronous metastases develop in approximately one third of all prostate cancer (PCa) patients. Main sites of metastasis include lymph nodes, bone, lung and liver. Secondary peritoneal carcinomatosis is very rare in PCa, with only a few published cases. ⁶⁸Ga-PSMA PET–based imaging is a promising tool for staging and restaging PCa. We report a case of ⁶⁸Ga-PSMA PET/MR–positive peritoneal metastasis as site of primary relapse after definitive PCa treatment in a 58-year-old man with a prostate-specific antigen of 30 ng/mL at time of the study. Exploratory laparoscopy and subsequent histopathologic examination confirmed nonascitic peritoneal PCa carcinomatosis.

Key Words: ⁶⁸Ga-PSMA, peritoneal carcinomatosis, PET/MRI, prostate cancer

(*Clin Nucl Med* 2017;00: 00–00)

Received for publication March 8, 2017; revision accepted April 3, 2017.

From the *Department of Urology, University Hospital Zurich, University of Zurich, Zurich; †Department of Radiation Oncology, University Hospital Geneva, Geneva; and ‡Department of Nuclear Medicine, University Hospital Zurich, University of Zurich, Zurich, Switzerland.

Conflicts of interest and sources of funding: none declared.

Correspondence to: Irene A. Burger, MD, Department of Nuclear Medicine, University Hospital Zurich, Rämistrasse 100, 8091 Zürich, Switzerland. E-mail: Irene.burger@usz.ch.

Copyright © 2017 Wolters Kluwer Health, Inc. All rights reserved.

ISSN: 0363-9762/17/0000–0000

DOI: 10.1097/RLU.0000000000001703

REFERENCES

1. Rauscher I, Maurer T, Beer AJ, et al. Value of ⁶⁸Ga-PSMA HBED-CC PET for the assessment of lymph node metastases in prostate cancer patients with biochemical recurrence: comparison with histopathology after salvage lymphadenectomy. *J Nucl Med*. 2016;57:1713–1719.
2. Kabasakal L, Demirci E, Nematyazar J, et al. The role of PSMA PET/CT imaging in restaging of prostate cancer patients with low prostate-specific antigen levels. *Nucl Med Commun*. 2017;38:149–155.
3. Eiber M, Maurer T, Souvatzoglou M, et al. Evaluation of hybrid ⁶⁸Ga-PSMA ligand PET/CT in 248 patients with biochemical recurrence after radical prostatectomy. *J Nucl Med*. 2015;56:668–674.
4. Freitag MT, Radtke JP, Hadaschik BA, et al. Comparison of hybrid (68)Ga-PSMA PET/MRI and (68)Ga-PSMA PET/CT in the evaluation of lymph node and bone metastases of prostate cancer. *Eur J Nucl Med Mol Imaging*. 2016;43:70–83.
5. Taneja S, Taneja R, Kashyap V, et al. ⁶⁸Ga-PSMA uptake in hepatocellular carcinoma. *Clin Nucl Med*. 2017;42:e69–e70.
6. Taywade SK, Damle NA, Bal C. PSMA expression in papillary thyroid carcinoma: opening a new horizon in management of thyroid cancer? *Clin Nucl Med*. 2016;41:e263–e265.
7. Shetty D, Loh H, Bui C, et al. Elevated ⁶⁸Ga prostate-specific membrane antigen activity in metastatic non–small cell lung cancer. *Clin Nucl Med*. 2016;41:414–416.
8. Nomura N, Pastorino S, Jiang P, et al. Prostate specific membrane antigen (PSMA) expression in primary gliomas and breast cancer brain metastases. *Cancer Cell Int*. 2014;14:26.
9. Henninger M, Maurer T, Hacker C, et al. ⁶⁸Ga-PSMA PET/MR showing intense PSMA uptake in nodular fasciitis mimicking prostate cancer metastasis. *Clin Nucl Med*. 2016;41:e443–e444.
10. Sheng J, Findley TW, Sadeghi-Nejad H. Isolated non-ascitic peritoneal carcinomatosis from metastatic prostate cancer. *Urol Case Rep*. 2016;10:14–15.
11. Sridhar S, Nguyen BD. Education and Imaging. Gastrointestinal: positron emission tomography/computed tomography imaging of peritoneal carcinomatosis secondary to prostate adenocarcinoma. *J Gastroenterol Hepatol*. 2016;31:704.
12. Gungor S, Asa S, Kupik O. FDG PET/CT in peritoneal metastasis from prostate cancer. *Clin Nucl Med*. 2016;41:e424–e425.
13. Petrakis D, Pentheroudakis G, Kamina S, et al. An unusual presentation of a patient with advanced prostate cancer, massive ascites and peritoneal metastasis: case report and literature review. *J Adv Res*. 2015;6:517–521.




FIGURE 1. ^{68}Ga prostate-specific membrane antigen (PSMA) PET/MRI. A 58-year-old man previously treated with robotic-assisted radical prostatectomy and lymph node sampling 11 years ago for a localized high-risk prostate cancer (pT2c, pN0 [0/2], cM0, R0, Gleason 4 + 4 = 8, preoperative prostate-specific antigen [PSA] 40 ng/mL), who also received salvage radiation therapy combined with 12 months of androgen deprivation for a para-anastomotic recurrence 18 months after surgery, was referred for ^{68}Ga -PSMA PET/MR for restaging because of previous repeated negative PET imaging (^{18}F -choline and ^{68}Ga -bombesin) despite continuously rising PSA values (PSA at time of imaging 30 ng/mL, with an estimated PSA doubling time of 11 months). **A**, Coronal and sagittal MIP of the ^{68}Ga -PSMA PET/MR showed a single PSMA-positive hepatic lesion of 1.5×2.7 cm with a SUVmax of 25.9. There was no evidence for local recurrence in the prostate bed on **(B)** axial T2-weighted images or **(C)** axial-fused PET/MR. No positive lymph nodes, bone metastases, or any sign of ascites was detected. On the axial **(D)** fat and **(E)** water weighted T1 3-dimensional dual-echo gradient-echo pulse DIXON-LAVA-Flex sequences, the lesion was slightly hypointense (arrow), with **(F)** high PSMA accumulation on axial PET and **(G)** fused PET/MR.




FIGURE 2. Primovist liver MRI. Because an intrahepatic lesion could not be ruled out on the PSMA PET/MR, the patient underwent a dedicated liver MRI to further characterize the PSMA-positive lesion. **A**, The Primovist MRI revealed a single peritoneal implant located on the falciform ligament without contrast enhancement (yellow arrow, T1 FFE with fat saturation, 20 minutes after contrast injection). **B**, On axial diffusion-weighted images ($b = 800$), a clear diffusion restriction was documented (arrow), further suggesting a peritoneal metastasis. Exploratory laparoscopy confirmed the lesion and revealed a disseminated peritoneal carcinomatosis. The histopathologic analyses of the biopsy specimens of the falciform ligament lesion and several other peritoneal implants of 1 to 2 mm confirmed an infiltration by an undifferentiated prostate adenocarcinoma Gleason 4 + 5 = 9. ^{68}Ga -PSMA PET imaging offers promising possibilities in restaging of biochemical recurrence, especially at very low PSA values of less than 1 ng/mL.^{1–3} In addition, an accurate detection of nodal and bone metastases has been demonstrated.⁴ Besides primary prostate cancer and its metastases, ^{68}Ga -PSMA uptake has been reported in hepatocellular carcinoma and different other malignant and nonmalignant entities with high neovascularity.^{5–9} Peritoneal carcinomatosis in metastatic prostate cancer is extremely rare and has been described in only few single case reports.^{10–13} ^{68}Ga -PSMA-based imaging might offer improved detection rates for such occult visceral metastases.